

## Monochromatic Trees with Respect to Edge Partitions\*

VOJTĚCH RÖDL

*Emory University, Department of Mathematics,  
Atlanta, Georgia 30322*

AND

BERND VOIGT

*Fakultät für Mathematik, Universität Bielefeld,  
4800 Bielefeld 1, Germany*

Received August 7, 1984

It is shown, that for every infinite cardinal  $\kappa$  there exists a graph  $F$  on  $\kappa$  vertices satisfying  $F \rightarrow (T)_{\lambda}^{\text{edges}}$  for every tree  $T$  on  $\kappa$  vertices and all  $\lambda$  satisfying  $\text{cf } \kappa \rightarrow (\omega)_{\lambda}^3$ . © 1993 Academic Press, Inc.

### 1. PARTITIONING EDGES

Erdős, Hajnal, and Pósa asked, whether for every (infinite) graph  $G$  there exists a graph  $F$ , such that for every two-coloring of the edges of  $F$  there is some monochromatic and induced copy of  $G$  in  $F$ . For finite (resp. countable) graphs  $G$  the answer is known to be affirmative, cf., [2, 3, 7].

With respect to uncountable graphs the results turn out to depend very much on the underlying set theoretical assumptions, compare [8, 9].

In this paper we investigate the particular case, when  $G$  is a tree on  $\kappa$  vertices. Our results do not use set theoretic assumptions and in most cases they give the smallest possible size of the Ramsey graphs.

*Notations.* For graphs  $F$  and  $H$  we denote by  $(F_H)$  the set of all induced subgraphs of  $F$  which are isomorphic to  $H$ . " $F \rightarrow (G)_{\lambda}^H$ ," where  $\lambda$  is some cardinal, abbreviates the following statement: *for every  $\lambda$ -coloring  $\Delta: (F_H) \rightarrow \lambda$  there exists some induced  $G$ -subgraph  $G^* \in (F_G)$  such that  $\Delta|_{(G^*_H)}$  is a constant mapping.* In particular, if  $H$  is the complete graph on two

\* With revisions from June 1991.

vertices (i.e.,  $H$  is a single edge), we write  $F \rightarrow (G)_\lambda^c$ . We say that  $H$  is weak subgraph of  $H$  if a subgraph which is not necessarily induced. The weak Ramsey-arrow " $F \rightarrow (G)_\lambda^{c''}$ " is defined as above, however, just referring to weak subgraphs. If  $H$  is the complete graph on two vertices (i.e.,  $H$  is a single edge), we write  $F \rightarrow (G)_\lambda^c$ . For infinite cardinals  $\kappa$ , the  $\kappa$ -regular tree, i.e., every vertex has degree  $\kappa$ , is denoted by  $T_\kappa$ .

The main result of our paper is the following:

**THEOREM A.** *For every infinite cardinal  $\kappa$  there exists a graph  $G_\kappa$  on  $\kappa$  vertices, such that  $G_\kappa \rightarrow (T_\kappa)_\lambda^c$  holds for all cardinals  $\lambda$  satisfying  $\text{cf}(\kappa) \rightarrow (\omega)_\lambda^3$ .*

From Ramsey's theorem we have the following corollary:

**COROLLARY A1.** *For every infinite cardinal  $\kappa$  there exists a graph  $G_\kappa$  on  $\kappa$  vertices, such that  $G_\kappa \rightarrow (T_\kappa)_n^c$  holds for every positive integer  $n$ .*

This improves the result of [6], where it has been shown that there exists a graph  $F$  on  $\exp(\kappa)$  vertices satisfying  $F \rightarrow (T_\kappa)_n^c$  for every positive integer. As by [4],  $(2^{2'})^+ \rightarrow (\omega)_\lambda^3$  for every infinite cardinal  $\lambda$ ; hence we have another corollary:

**COROLLARY A2.** *For every infinite cardinal  $\lambda$  there exists a graph  $G_\kappa$  on  $\kappa = (2^{2'})^+$  vertices satisfying  $G_\kappa \rightarrow (T_\kappa)_\lambda^c$ .*

*Proof of Theorem A.* Let  $\kappa$  be an infinite cardinal. The shift graph  $G_\kappa = (V_\kappa, E_\kappa)$  on  $\kappa$  is defined as follows:  $V_\kappa = [\kappa]^2$ ,  $E_\kappa = \{\{A, B\} \in [V_\kappa]^2 \mid \max A = \min B\}$ . Thus every three-element subset  $\{x, y, z\} \in [\kappa]^3$ , say  $x < y < z$ , corresponds to the edge  $\{\{x, y\}, \{y, z\}\} \in E_\kappa$ , and vice versa. We shall show that  $G_\kappa$  has the desired properties.

**CLAIM 1.** *Let  $W \subseteq V_\kappa$  be a set of vertices and let  $E^* \subseteq E_\kappa \cap [W]^2$  be a set of edges of  $G_\kappa$ , such that the graph  $(W, E^*)$  has minimal degree  $\kappa$ . Then  $(W, E^*)$  contains an induced copy of  $T_\kappa$ , which also is an induced subgraph with respect to  $G_\kappa$ .*

*Proof of Claim 1.* Using transfinite recursion, we shall show that there exists a one-to-one mapping  $\varphi: V(T_\kappa) \rightarrow W$  such that for all  $x, y \in V(T_\kappa)$  holds (i)  $\{x, y\} \in E(T_\kappa)$  implies that  $\{\varphi(x), \varphi(y)\} \in E^*$  and (ii)  $\{x, y\} \notin E(T_\kappa)$  implies that  $\{\varphi(x), \varphi(y)\} \notin E_\kappa$ . Then  $\{\varphi(x) \mid x \in V(T_\kappa)\}$  is the desired copy of  $T_\kappa$ .

Let  $\leq$  be a well-ordering of  $V(T_\kappa)$  having order type  $\kappa$  and let  $r \in V(T_\kappa)$  be its least element. Put  $R_0 = \{r\}$  and define  $\varphi_0: R_0 \rightarrow W$  arbitrarily.

Assume, by induction, that for every  $v < \xi < \kappa$  sets  $R_v \subseteq V(T_\kappa)$  and mappings  $\varphi_v: R_v \rightarrow W$  have been defined such that

- (1)  $R_v \subseteq R_\mu$  and  $\varphi_\mu \upharpoonright R_v = \varphi_v$ , for all  $v < \mu < \xi$ ,
- (2) for all  $v < \mu < \xi$  and all  $x \in R_v$  and  $y \in R_\mu$  we have:
  - (2i)  $\{x, y\} \in E(T_\kappa)$  implies that  $\{\varphi_v(x), \varphi_\mu(y)\} \in E^*$ ,
  - (2ii)  $\{x, y\} \notin E(T_\kappa)$  implies that  $\{\varphi_v(x), \varphi_\mu(y)\} \notin E_\kappa$ ,
  - (2iii)  $\max \varphi_v(x) \neq \max \varphi_\mu(y)$ ,
- (3) every  $R_v$  is a connected subset of vertices of  $T_\kappa$  and  $|R_v| \leq |v| + 1$ .

Let  $z \in V(T_\kappa)$  be the least element (w.r.t.  $\leq$ ) which is not already contained in  $\bigcup_{v < \xi} R_v$ , but which is connected by an edge in  $E(T_\kappa)$  to some (uniquely determined, as  $T_\kappa$  is a tree) vertex  $x \in \bigcup_{v < \xi} R_v$ . Say  $x \in R_\mu$  and  $\varphi_\mu(x) = \{\alpha, \beta\}$ . Since  $|\bigcup_{v < \xi} R_v| \leq |\xi| + 1 < \kappa$  and as, with respect to  $E^*$ , vertex  $\{\alpha, \beta\} \in W$  has degree  $\kappa$ , there exists some  $\gamma < \kappa$  such that  $\{\{\alpha, \beta\}\{\beta, \gamma\}\} \in E^*$ , but  $\gamma \notin \varphi_v(y)$  for all  $v < \xi$  and  $y \in R_v$ .

Put  $R_\xi = \{z\} \cup \bigcup_{v < \xi} R_v$  and  $\varphi_\xi(z) = \{\beta, \gamma\}$  and  $\varphi_\xi(y) = \varphi_v(y)$  for all  $v < \xi$  and  $y \in R_v$ . Then again properties (1), (2), and (3) are satisfied, and, by construction,  $\varphi = \bigcup_{v < \kappa} \varphi_v$  has the desired properties. ■

**CLAIM 2.** Let  $\lambda$  satisfy  $\text{cf} \kappa \rightarrow (\omega)_\lambda^3$  and let  $\Delta: E_\kappa \rightarrow \lambda$  be a mapping. Then there exists a  $\kappa$ -subset  $Y \in [\kappa]^\kappa$  and there exists some  $\eta < \lambda$ , such that the graph  $(Y, E^{**})$ , where  $E^{**} = \{\{A, B\} \in [Y]^2 \cap E_\kappa \mid \Delta(\{A, B\}) = \eta\}$ , is  $\kappa$ -regular.

*Proof of Claim 2.* Assume to the contrary that Claim 2 fails to be true. Then, by transfinite induction, one easily defines for every  $\xi < \lambda$  a well-ordering  $<_\eta$  of  $[\kappa]^2$  with the following property:

$$|\{B \in [\kappa]^2 \mid A <_\xi B, \{A, B\} \in E_\kappa \text{ and } \Delta(\{A, B\}) = \eta\}| < \kappa$$

for every  $A \in [\kappa]^2$ . (\*)

Consider the following recursive construction: Put  $\alpha_0 = 0$  and  $R_0 = \kappa$  and assume, by induction, that for every  $v < \xi < \text{cf} \kappa$  ordinals  $\alpha_v < \kappa$  and subsets  $R_v \subseteq \kappa$  have been defined such that (i)  $|\kappa \setminus R_v| < \kappa$  and (ii)  $\alpha_v = \min \bigcap_{\mu < v} R_\mu$  for every  $v < \xi$ .

Put  $R'_\xi = \bigcap_{v < \xi} (R_v \setminus \{\alpha_v\})$ . Note that  $|\kappa \setminus R'_\xi| < \kappa$ , as  $\xi < \text{cf} \kappa$ . Put  $\alpha_\xi = \min R'_\xi$ . For every ordinal  $v < \xi$  consider the vertex  $A = \{\alpha_v, \alpha_\xi\}$  and remove from  $R'_\xi$  all ordinals  $\beta$  with  $A <_\eta \{\alpha_\xi, \beta\}$ ,  $\{A, \{\alpha_\xi, \beta\}\} \in E_\kappa$ , and  $\Delta(\{A, \{\alpha_\xi, \beta\}\}) = \eta$  for some  $\eta < \lambda$ . According to (\*), the remaining subset, call it  $R_\xi$ , satisfies (i), and, moreover,

$$\{\alpha_\xi, \gamma\} <_\eta \{\alpha_v, \alpha_\xi\} \quad \text{for every } \gamma \in R_\xi \text{ and every } v < \xi$$

with  $\Delta(\{\{\alpha_v, \alpha_\xi\}, \{\alpha_\xi, \gamma\}\}) = \eta$ . (\*\*)

Consider the set  $\mathcal{A} = \{\alpha_\xi \mid \xi < \text{cf } \kappa\}$ . As  $\text{cf } \kappa \rightarrow (\omega)_\lambda^3$ , there exists an infinite subset  $\{\beta_0, \beta_1, \dots\} \subseteq \mathcal{A}$ , say,  $\beta_0 < \beta_1 < \dots$ , and there exists some  $\eta < \lambda$  such that

$$\mathcal{A}(\{\{\beta_i, \beta_{i+1}\}, \{\beta_{i+1}, \beta_{i+2}\}\}) = \eta \quad \text{for every } i < \omega.$$

By (\*\*), this implies that

$$\{\beta_{i+1}, \beta_{i+2}\} <_\eta \{\beta_i, \beta_{i+1}\} \quad \text{for every } i < \omega,$$

contradicting that  $<_\xi$  is a well-ordering. Thus the claim is proved. ■

Finally, Theorem A is an immediate conclusion drawn from Claim 2 and Claim 1. ■

*Remark 1.* For later use let us note that a straightforward modification of the proof of Claim 1 establishes the following lemma:

**LEMMA 1.** *Let  $G = (V, E)$  be a graph of minimal degree  $\kappa$ , where  $\kappa$  is an infinite cardinal. Then  $G$  contains a (non-induced) subgraph isomorphic to  $T_\kappa$ .*

*Proof.* Again, let  $\leq$  be a well-ordering of  $V(T_\kappa)$  having order type  $\kappa$  and let  $r$  be its minimal element. Put  $R_0 = \{r\}$  and define  $\varphi_0: R_0 \rightarrow V$  arbitrarily. Assume that for every  $v < \xi < \kappa$  connected subsets  $R_v \subseteq V(T_\kappa)$  and embeddings  $\varphi_v: R_v \rightarrow V$  have been defined such that  $R_v \subseteq R_\mu$  and  $\varphi_\mu \upharpoonright R_v = \varphi_v$  for all  $v < \mu < \xi$ . Let  $z \in V(T_\kappa)$  be the least element of  $V(T_\kappa) \setminus \bigcup_{v < \xi} R_v$  which is connected by an edge to some  $x \in \bigcup_{v < \xi} R_v$ . Say  $x \in R_\mu$  and  $\varphi_\mu(x) = a$ . As  $a$  is joined by an edge to at least  $\kappa$  many vertices in  $V$ , there exists some  $b \in V \setminus \bigcup_{v < \xi} \varphi_v(R_v)$  such that  $\{a, b\} \in E$ . Put  $R_\xi = \{z\} \cup \bigcup_{v < \xi} R_v$  and define  $\varphi_\xi(z) = b$  and  $\varphi_\xi(y) = \varphi_v(y)$  for every  $y \in R_v$ ,  $v < \xi$ . By construction then  $\varphi = \bigcup_{v < \kappa} \varphi_v$  is a weak embedding of  $T_\kappa$  into  $G$ . ■

With a different proof, Lemma 1 can be found in [1].

*Remark 2.* The condition “ $\text{cf } \kappa \rightarrow (\omega)_\lambda^3$ ” cannot be weakened to “ $\kappa \rightarrow (\omega)_\lambda^3$ .” This can be seen, e.g., by considering  $K_{\aleph_\omega}$ , the complete graph on  $\aleph_\omega$  vertices:

**LEMMA 2.** *Consider  $K_{\aleph_\omega} = (\aleph_\omega, [\aleph_\omega]^2)$ , the complete graph on  $\aleph_\omega$  vertices. Then  $K_{\aleph_\omega} \not\rightarrow^{weak} (T_{\aleph_\omega})_\omega^e$ ; i.e., there exists a mapping  $\Delta: [\aleph_\omega]^2 \rightarrow \omega$  such that no (weak) subgraph  $(\aleph_\omega, \Delta^{-1}(\xi))$ ,  $\xi < \omega$ , contains some  $\aleph_\omega$ -regular (weak) subgraph, where “weak” means “not necessarily induced.”*

*Proof.* Put  $\Delta(\{\alpha, \beta\}) = (i, k)$  for all  $\aleph_{i-1} \leq \alpha < \aleph_i$  and  $\aleph_{k-1} \leq \beta < \aleph_k$ , where  $\aleph_{-1} = 0$  for convenience. ■

*Remark 3.* Corollary A2 suggests considering the function  $f$ , which is defined for infinite cardinals  $\lambda$  such that  $f(\lambda)$  is the least cardinal  $\kappa$  for which there exists a graph  $G$  on  $\kappa$  vertices with  $G \rightarrow (T_\kappa)_\lambda^e$ . From Corollary A2 it follows that  $f(\lambda) \leq (2^{2^\lambda})^+$ . On the other hand, we can show that  $\exp(\lambda)^+ \leq f(\lambda)$ .

**LEMMA 3.** *For  $\kappa = \exp(\lambda)$  the complete graph on  $\kappa$  vertices,  $K_\kappa$ , satisfies  $K_\kappa \not\rightarrow_{\text{weak}} (T_\kappa)_\lambda^e$ ; i.e., there exists a mapping  $\Delta: [\exp(\lambda)]^2 \rightarrow \lambda$  such that no (weak) subgraph  $(\exp(\lambda), \Delta^{-1}(\xi))$ ,  $\xi < \lambda$ , contains some  $\exp(\lambda)$ -regular subgraph.*

*Proof.* Let us view  $\exp(\lambda)$  as the set of 0-1 sequences of length  $\lambda$ . Let  $\leq^*$  be a well-ordering of type  $\exp(\lambda)$  of these sequences. The mapping  $\Delta$  is defined as follows:

$$\begin{aligned} \Delta(\{(x_i)_{i < \lambda}, (y_i)_{i < \lambda}\}) &= (\xi, 0) && \text{if } \min\{i < \lambda \mid x_i \neq y_i\} = \xi \\ &&& \text{and } x_\xi = 0 \text{ and } (x_i)_{i < \lambda} <^* (y_i)_{i < \lambda} \\ &= (\xi, 1) && \text{if } \min\{i < \lambda \mid x_i \neq y_i\} = \xi \\ &&& \text{and } x_\xi = 0 \text{ and } (x_i)_{i < \lambda} >^* (y_i)_{i < \lambda}. \end{aligned}$$

Intuitively, we did the following: The complete graph  $[\exp(\lambda)]^2$  is partitioned into  $\lambda$  many bipartite graphs, where  $\{(x_i)_{i < \lambda}, (y_i)_{i < \lambda}\}$  belongs to the  $\xi$ th bipartite graph iff  $\xi = \min\{i \mid x_i \neq y_i\}$ . For each such bipartite graph we have a left-hand side (those  $(x_i)_{i < \lambda}$  with  $x_\xi = 0$ ) and a right-hand side (those  $(x_i)_{i < \lambda}$  with  $x_\xi = 1$ ). Now the edge  $\{(x_i)_{i < \lambda}, (y_i)_{i < \lambda}\}$  belongs to class  $(\xi, 0)$  iff  $(x_i)_{i < \lambda} <^* (y_i)_{i < \lambda}$ . ■

We conjecture that  $f(\lambda) = \exp(\lambda)^+$ . But we can only show the following:

**LEMMA 4.**  $K_{\exp(\lambda)^+} \not\rightarrow_{\text{weak}} (T_{\exp(\lambda)^+})_\lambda^e$  for every infinite cardinal  $\lambda$ .

*Proof.* In view of Lemma 1 it suffices to prove the following claim:

**CLAIM 3.** *For every mapping  $\Delta: [\exp(\lambda)^+]^2 \rightarrow \lambda$  there exists a subset  $Y \in [\exp(\lambda)^+]^{\exp(\lambda)^+}$  and there exists some  $\eta < \lambda$ , such that for every  $\alpha \in Y$  there exists  $Z_\alpha \in [Y]^{\exp(\lambda)^+}$  with  $\Delta(\{\alpha, \beta\}) = \eta$  for every  $\beta \in Z_\alpha$ .*

*Proof of Claim 3.* We use a slight modification of the proof of Claim 2. Assume to the contrary, that the claim fails to be true. Then for every  $\xi < \lambda$  there exists a well-ordering  $<_\eta$  of  $\exp(\lambda)^+$  such that

$$\begin{aligned} |\{\beta \in \exp(\lambda)^+ \mid \alpha <_\eta \beta \text{ and } \Delta(\{\alpha, \beta\}) = \eta\}| \\ < \exp(\lambda)^+ \quad \text{for every } \alpha \in \exp(\lambda)^+. \end{aligned} \quad (*)$$

Let  $\xi < (\exp \lambda)^+$  and assume by induction that for every  $v < \xi$  ordinals  $\alpha_v$  and sets  $R_v \subseteq (\exp \lambda)^+$  have been defined such that (i)  $|(\exp \lambda)^+ - R_v| < (\exp \lambda)^+$  and  $\alpha_\xi = \min \bigcap_{v < \xi} R_v$  for every  $v < \xi$ . Put  $\alpha_\xi = \min \bigcap_{v < \xi} R_v$ , where, for convenience,  $\bigcap_{\emptyset} R_v = (\exp \lambda)^+$ . For every ordinal  $v \leq \xi$  remove from  $\bigcap_{v < \xi} R_v$  all ordinals  $\beta$  with  $\alpha_v <_\eta \beta$  and  $\Delta(\{\alpha_v, \beta\}) = \eta$ . According to (\*) and (i), the remaining set, call it  $R_\xi$ , again satisfies (i). Moreover,

$$\beta <_\eta \alpha_v \text{ for every } v \leq \xi \text{ and every } \beta \in R_\xi \text{ with } \Delta(\{\alpha_v, \beta\}) = \eta. \quad (**)$$

Consider the set  $\mathcal{A} = \{\alpha_v \mid v < (\exp \lambda)^+\}$ . As  $(\exp \lambda)^+ \rightarrow (\omega)_\lambda^2 [4]$ , there exists an infinite subset  $\{\beta_0, \beta_1, \dots\} \subseteq \mathcal{A}$ , say  $\beta_0 < \beta_1 < \dots$ , and there exists  $\eta < \lambda$  such that  $\Delta(\{\beta_i, \beta_{i+1}\}) = \xi$  for every  $i < \omega$ . By (\*\*) this implies that  $\beta_{i+1} <_\eta \beta_i$  for every  $i < \omega$ , contradicting that  $<_\eta$  is a well-ordering. ■

## 2. RAMIFICATIONS

### 1. Partitioning Vertices

If  $H$  is a single vertex, we write  $F \rightarrow (G)_\lambda^c$  instead of  $F \rightarrow (G)_\lambda^H$ . Using the same method as for Theorem A we can show:

**THEOREM B.** *For every infinite cardinal  $\kappa$  there exists a graph  $G_\kappa$  on  $\kappa$  vertices, such that  $G_\kappa \rightarrow (T_\kappa)_\lambda^c$  for all satisfying  $\text{cf} \kappa \rightarrow (\omega)_\lambda^2$ .*

*Proof of Theorem B.* We use the same graph  $G_\kappa$  as for Theorem A. Let  $\Delta: [\kappa]^2 \rightarrow \lambda$  be an arbitrary mapping. This can be viewed as an edge-coloring of the complete graph  $K_\kappa$ . Hence, by Lemma 4, there exists a weakly embedded copy of  $T_\kappa$  in  $K_\kappa$ , say with edges  $W \subseteq [\kappa]^2$ , which is colored monochromatically. Viewing these edges as vertices of  $G_\kappa$ , shows that  $(W, [W]^2 \cap E_\kappa)$  has minimal degree  $\kappa$ . Hence, by Claim 1, Theorem B follows. ■

*Remark.* Theorem B suggests considering the function  $g(\lambda)$  which is defined for infinite cardinals such that  $g(\lambda)$  is the minimal cardinal  $\kappa$  such that there exists a graph  $G$  on  $\kappa$  vertices satisfying  $G \rightarrow (T_\kappa)_\lambda^c$ . Clearly,  $\lambda^+ \leq g(\lambda)$ , and from Theorem B we infer that  $g(\lambda) \leq (\exp \lambda)^+$ . Again, we conjecture that the lower bound gives the right value.

## 3. DIRECTED GRAPHS

Let us denote by  $\tilde{T}_\kappa$  the directed  $\kappa$ -regular arborescence; i.e., call some distinguished vertex  $r \in V(T_\kappa)$  the root of  $\tilde{T}_\kappa$  and direct all edges of  $E(T_\kappa)$  in such a way that every vertex  $x \in V(T_\kappa) \setminus \{r\}$  can be reached from  $r$  by a

directed path. By  $\vec{T}_\kappa$  we denote the directed tree obtained from  $\tilde{T}_\kappa$  by reversing the orientation.

The directed version of Theorem A then says:

**THEOREM C.** *For every infinite cardinal  $\kappa$  there exists a directed graph  $\vec{G}_\kappa$  on  $\kappa$  vertices, such that  $\vec{G} \rightarrow (\vec{T}_\kappa)_\lambda^c$  holds for all cardinals  $\lambda$  satisfying  $\text{cf } \kappa \rightarrow (\omega)_\lambda^3$ .*

**COROLLARY C1.** *For every infinite cardinal  $\kappa$  there exists a directed graph  $\vec{G}_\kappa$  on  $\kappa$  vertices, such that  $\vec{G}_\kappa \rightarrow (\vec{T}_\kappa)_n^c$  holds for every positive integer  $n$ .*

**COROLLARY C2.** *For every infinite cardinal  $\kappa$  there exists an undirected graph  $G_\kappa$  on  $\kappa$  vertices, such that for every orientation  $\vec{G}_\kappa^*$  of  $G_\kappa$ ,  $\vec{G}_\kappa^*$  contains some induced copy of  $\vec{T}_\kappa$  or it contains some induced copy of  $\vec{T}_\kappa$ .*

*Proof of Corollary C2.* Let  $G_\kappa$  be the undirected version of the graph  $\vec{G}_\kappa$  from Corollary C1. Given an arbitrary orientation  $\vec{G}_\kappa^*$ , impose a two-coloring on the edges of  $\vec{G}_\kappa$  by comparing the two orientations and apply Corollary 1. ■

Corollaries C1 and C2 improve results of [6], where the existence of such graphs  $\vec{G}_\kappa$  on  $\exp(\kappa)$  vertices has been established.

**COROLLARY C3.** *For every infinite cardinal  $\lambda$  there exists a directed graph  $\vec{G}_\kappa$  on  $\kappa$  vertices, where  $\kappa = \exp_2(\lambda)^+$ , such that  $\vec{G}_\kappa \rightarrow (\vec{T}_\kappa)_\lambda^c$ .*

*Proof of Theorem C.* Let  $G_\kappa$  be the graph from Theorem A. Direct each edge  $\{\{\alpha, \beta\}, \{\beta, \gamma\}\}$  from  $\{\alpha, \beta\}$  to  $\{\beta, \gamma\}$ . Then the proof of Theorem A actually establishes Theorem C. ■

## REFERENCES

1. L. D. ANDERSON AND C. THOMASSEN, The cover index of finite graphs, *Aequationes Math.* **20** (1980), 244–251.
2. W. DEUBER, A Generalization of Ramsey's theorem, in "Infinite and Finite Sets," Vol. I (A. Hajnal, R. Rado, and V. T. Sós, Eds.), Colloq. Math. Soc. Bolyai, Vol. 10, pp. 323–332, North-Holland, Amsterdam, 1975.
3. P. ERDŐS, A. HAJNAL, AND L. PÓSA, Strong embeddings of graphs into colored graphs, in "Infinite and Finite Sets" (A. Hajnal, R. Rado, and V. Sós, Eds.), Colloq. Math. János Bolyai, Vol. 10, pp. 584–596, North-Holland, Amsterdam, 1975.
4. P. ERDŐS AND R. RADO, A partition calculus in set theory, *Bull. Amer. Math. Soc.* **62** (1950), 427–489.
5. R. L. GRAHAM, B. L. ROTHCHILD, AND J. H. SPENCER, "Ramsey Theory," Wiley, New York, 1980.

6. H. J. PRÖMEL, C. THOMASSEN, AND B. VOIGT, Induced oriented trees in graphs, unpublished manuscript, Bielefeld, 1980.
7. V. RÖDL, "Generalization of Ramsey's Theorem and Dimension of Graphs," Charles University Prague, Thesis 1973 [Czech]; English translation, Generalization of Ramsey's theorem, in "Graphs Hypergraphs and Block Systems," pp. 211–220, Zielona Gora, 1976.
8. A. HAJNAL AND P. KOMJÁTH, Embedding graphs into colored graphs, *Trans. Amer. Math. Soc.* **307** (1988), 395–409.
9. S. SHELAH, Consistency of positive partition theorems for graphs and models, in "Set Theory and Its Applications" (J. Steprāns and S. Watson, Eds.), pp. 167–193, Springer-Verlag, New York/Berlin, 1989.